

United States Department of Agriculture – Agricultural Research Service research on alternatives to methyl bromide: pre-plant and post-harvest^{†‡}

Sally M Schneider,^{1*} Erin N Roskopf,² James G Leesch,¹ Daniel O Chellemi,² Carolee T Bull³ and Mark Mazzola⁴

¹USDA-ARS, San Joaquin Valley Agricultural Sciences Center, Parlier, CA 93648, USA

²USDA-ARS, U.S. Horticultural Research Laboratory, Ft. Pierce, FL 34945, USA

³USDA-ARS, Crop Improvement and Protection Research Unit, Salinas, CA 93905, USA

⁴USDA-ARS, Tree Fruit Research Laboratory, Wenatchee, WA 98801, USA

Abstract: Methyl bromide is a widely used fumigant for both pre-plant and post-harvest pest and pathogen control. The Montreal Protocol and the US Clean Air Act mandate a phase-out of the import and manufacture of methyl bromide, beginning in 2001 and culminating with a complete ban, except for quarantine and certain pre-shipment uses and exempted critical uses, in January 2005. In 1995, ARS built on its existing programs in soil-borne plant pathology and post-harvest entomology and plant pathology to initiate a national research program to develop alternatives to methyl bromide. The focus has been on strawberry, pepper, tomato, perennial and nursery cropping systems for pre-plant methyl bromide use and fresh and durable commodities for post-harvest use. Recently the program has been expanded to include research on alternatives for the ornamental and cut flower cropping systems. An overview of the national research program is presented. Results from four specific research trials are presented, ranging from organic to conventional systems. Good progress on short-term alternatives is being made. These will be used as the foundation of integrated management systems which begin with pre-plant management decisions and continue through post-harvest processing.

Published in 2003 for SCI by John Wiley & Sons, Ltd.

Keywords: methyl bromide; pre-plant; post-harvest; strawberry; tomato; pepper; flower; ornamental; apple; grape; organic farming; chemical control; cultural control; biological control; plant pathogens; nematodes; weeds

1 INTRODUCTION

Methyl bromide is a critical element in pre-plant management of soil-borne pests and pathogens in high value fruits, nuts, vegetables, nursery and ornamental crops, and in post-harvest management of pests and pathogens on fresh produce and durable commodities. The Montreal Protocol, an international treaty, and the US Clean Air Act restricted availability of methyl bromide; beginning in January 2001, its use was restricted to 50% of the amount used in the baseline year of 1991, further restricted to 30% of the baseline in 2003 and completely banned in 2005. Quarantine use of methyl bromide is exempted from the impending ban. US growers of high value crops are in dire need of alternative management strategies for both pre-plant and post-harvest uses. The availability of acceptable alternatives will impact upon

the supply and quality of these foods to both American consumers and the export market. In response to the urgent need, the United States Department of Agriculture—Agricultural Research Service (ARS) initiated a diversified research program in 1995 that spans 10 states and Washington, DC, expenditure for which has grown to approximately \$15 million in 2001. This paper overviews the research program and presents a sampling of research highlights.

2 PRE-PLANT

Pre-plant soil fumigation with methyl bromide is used to control 'replant disorder', weeds, and soil-borne pathogens for many high value crops including strawberries, tomatoes, peppers, ornamentals, nursery crops, grapes, tree fruit and nut trees. The phasing out

* Correspondence to: Sally M Schneider, Research Plant Pathologist, USDA-ARS, San Joaquin Valley Agricultural Services Center, 9611 So Riverbend Ave, Parlier, CA 93648, USA

E-mail: sschneider@fresno.ars.usda.gov

[†]One of a collection of papers on various aspects of agrochemicals research contributed by staff of the Agricultural Research Service of the United States Department of Agriculture, and collected and organized by Drs RD Wauchope, NN Ragsdale and SO Duke

[‡]This article is a US Government work and is in the public domain in the USA

(Received 1 November 2002; revised version received 11 February 2003; accepted 26 February 2003)

of methyl bromide raises two major issues. The first is to quickly find effective, alternative control measures. Methyl bromide can be used effectively against a broad spectrum of soil pests over a range of soil types, temperature and moistures resulting in greater flexibility and less risk of loss than is possible with other soil treatments. Unless a 'silver bullet' that is effective against a wide range of pests can be found, the first challenge will be to accurately diagnose the problem(s) in a specific field. Once the problem is identified, a management strategy must be generated that is (1) effective against the identified pest, (2) effective under the soil conditions found in that field, (3) economically feasible, and (4) environmentally acceptable. ARS is addressing this issue by testing plant growth promoting rhizobacteria (PGPR), suppressive soils, soil amendments, fallow, mulches, crop rotation, host resistance, new chemicals and new application technologies to deliver biological and chemical alternatives. The second issue is to increase understanding of the pathogens and soil factors limiting crop production. A long-term, integrated management approach requires a thorough understanding of biological, chemical and physical soil factors, their interactions and their spatial variability, and will include cultural, genetic, biological and chemical management strategies. The short-term solutions to methyl bromide alternatives are stepping stones to the long-term research into integrated systems.

2.1 Annual crops

Annual crops have traditionally relied on methyl bromide treatment before each cropping season to control soil-borne pathogens, nematodes and weeds. The level of pest and pathogen control achieved with methyl bromide in annual crops is often the cumulative result of annual fumigation over many years. As methyl bromide is phased out, unexpected pests and pathogens that were unknowingly controlled by the annual methyl bromide fumigation are likely to appear. ARS research has focused on strawberry and vegetable production systems in California and Florida. Recently, ARS created two new positions to address the needs of flower and ornamental production systems.

2.1.1 Florida tomato and pepper production

2.1.1.1 Overview

Florida is the leading producer of fresh market tomato and bell pepper in the USA, with 23 760 ha producing a crop valued at US \$745 million.¹ The fresh market tomato industry in Florida accounts for more than one-third of the US crop² and use of methyl bromide has been estimated to be as high as 94% of the acreage planted to tomatoes annually.³ Root-knot nematode, fungal plant pathogens and weeds are of great concern in the absence of methyl bromide. Together, the Florida tomato and pepper industries account for 8% of the global consumption of methyl bromide⁴ and, without viable alternatives, production is predicted to decline by 60% and 63% respectively.⁵ In Florida, growers use a 'raised-bed plastic mulch' production

system that consists of seedlings transplanted into pre-formed beds that have been fumigated with methyl bromide and covered with polyethylene plastic.^{6,7} Although there are registered soil fumigants that show promise for disease and nematode control, there are limited numbers of effective herbicide partners available for these crops.

2.1.1.2 Cultural and genetic control

Preliminary studies on the use of paper mulch for nutsedge control in the raised-bed vegetable production system have shown that paper provides excellent weed suppression that is equivalent to or better than the use of the combination of 1,3-dichloropropene (1,3-D; Telone) + chloropicrin and pebulate (Tillam) (Roskopf EN, unpublished). Nematode-susceptible bell pepper cultivars grown in fields previously cropped with the resistant cultivar Carolina Cayenne⁸ had reduced galling from the root-knot nematode, *Meloidogyne incognita* (Kof & White) Chitwood, and yield was 2.8 times greater than in plots previously cropped to a susceptible variety.⁹

2.1.1.3 Biological control

Research on the use of plant-growth-promoting-rhizobacteria (PGPR) in combination with soil disinfestation treatments contributed to the development of BioYield™. The combination of bacterial strains LS213 and LS256 with methyl bromide resulted in higher pepper yields than with methyl bromide alone.¹⁰ Fungal plant pathogens, developed as components of an integrated approach to weed management, are highly host specific and affect only the target weed.¹¹ *Dactylaria higginsii* (Luttrell) MB Ellis is being tested currently as an off-season treatment for control of nutsedge and as a post-emergence spray in combination with 1,3-D + chloropicrin for tomato production.¹² *Phomopsis amaranthicola* Roskopf, Charud., Shabana & Benny, a pigweed pathogen, is an excellent candidate for use in the ornamentals production system, where few herbicides are available.¹³ Non-pathogenic strains of *Fusarium oxysporum* Schlecht, isolated from tomato roots grown in a suppressive soil, provide significant and consistent control of Fusarium wilt of tomato.^{14,15}

2.1.1.4 Chemical control

On-going cooperative research projects with scientists at the University of Florida include various application techniques with existing chemical alternatives¹⁶ as well as the development of new chemistries, such as propargyl bromide¹⁷ and reduced-risk compounds.¹⁸ Several years of field trials have been conducted with Plant Pro45™, an iodine-based material, generating encouraging results in the control of root-knot nematode, the fungus *Fusarium oxysporum* f sp *basili* Tamietti & Matta, and several weed species.^{19–21}

2.2 Research highlight: large scale field trials

Large-scale field trials were instrumental in identifying technical problems not evident in small-scale research

plots, developing information on control of soil-borne pests under the range of environmental and cultural practices experienced by growers, generating information on costs incurred at the farm level, and providing growers with the experience to evaluate alternatives.

2.2.1 Soil solarization field trial

Initially, soil solarization was evaluated in Florida as a broadcast treatment and found to control several key soilborne pests.^{22–24} It was not compatible with local production systems, however, due to the increased cost of the plastic and the accumulation of storm water run-off during heavy rains. Soil solarization was adapted to local production systems by performing strip solarization with clear plastic on the same beds used for production. Solarization on raised beds produced higher soil temperatures than broadcast solarization on a flat surface, eliminating the border effect associated with soil solarization and improving its efficacy.²⁵ Painting the clear plastic with white paint allowed it to function as horticultural mulch and terminated the solarization period. In research trials, significant control of most soil-borne pests was obtained, the exception being rootknot nematode.^{25–27}

Twenty-one large-scale field trials of soil solarization have been conducted on a total of 44.1 ha over the years 1995–1999. To address the lack of control of root-knot nematode, some trials included fumigation with 1,3-D + chloropicrin and use of disease-resistant cultivars. Pest control and marketable yield were measured and compared with data obtained from adjacent methyl bromide treated areas. Weed control in solarized plots was comparable with that in adjacent methyl bromide plots in all locations except when purslane (*Portulaca oleracea* L) and Texas panicum (*Panicum texanum* Buckl) were present. Root gall ratings indicated that soil solarization did not

provide adequate control of root-knot nematodes. When solarization was combined with reduced rates of 1,3-dichloropropene + chloropicrin, reductions in root galling were similar to those achieved in adjacent methyl bromide fumigated plots. The incidence of soilborne diseases caused by *Fusarium oxysporum* f sp *lycopersici* Snyder & Hans, *Phytophthora capsici* Leonian and *Sclerotium rolfsii* Sacc were similar in solarized and methyl bromide treated plots in all trials except the 1999 tomato trial, where the incidence of *Fusarium* wilt was significantly higher in solarized beds. In the tomato field trials, marketable yields in solarized plots were between 85% and 106% of yields obtained in adjacent methyl bromide treated plots (Table 1). Soil solarization resulted in an average reduction in yield of 5% (Table 1). Fumigation with mixtures of 1,3-D and chloropicrin at the initiation of solarization did not improve yields, even though the control of nematodes was improved (Table 1). Large-scale field trials identified several technical problems not evident in smaller research plots. When using drip irrigation, the tube must be covered with soil during the solarization period to prevent melting. Paint coverage at the termination of the solarization period must be uniform and complete to prevent any additional solar radiation from penetrating the plastic and heating the soil to levels detrimental to the health of the transplants.

2.2.2 Broadcast fumigation field trial

When combined with the herbicide pebulate, shank injection of 1,3-D plus chloropicrin (Telone C17 and Telone C35) into beds prior to laying plastic has provided pest control and yields similar to those achieved with methyl bromide.^{28,29} Application of 1,3-D requires personal protective equipment (PPE) to be worn by workers during application.³⁰ If 1,3-D were applied via shank injection during bedding, which is the recommended procedure for methyl

Table 1. Marketable yields obtained from large-scale trials using soil solarization

Farm	Year	Crop	Treatment	Marketable yield (kg ha ⁻¹)	Comparison with methyl bromide (%)
1	1995	Tomato	Solarization	60 536	98
2	1995	Tomato	Solarization	54 320	106
3	1995	Tomato	Solarization	44 324	85
4	1996	Tomato	Solarization	48 244	95
5	1997	Tomato	Solarization	55 328	92
6	1998	Tomato	Solarization	36 512	96
Average				54 430	95
1	1995	Tomato	Solarization + 1,3-D:chloropicrin (83:17 mixture at 164 liter ha ⁻¹)	48 870	88
2	1996	Tomato	Solarization + 1,3-D:chloropicrin (83:17 mixture at 164 liter ha ⁻¹)	43 980	81
3	1996	Tomato	Solarization + 1,3-D (93 liter ha ⁻¹)	67 620	92
4	1999	Tomato	Solarization + 1,3-D (112 liter ha ⁻¹)	39 090	79
5	1999	Tomato	Solarization + 1,3-D:chloropicrin (65:35 at 327 liter ha ⁻¹)	45 030	94
Average				48 930	87

bromide application in Florida vegetable production, 30–50 workers would have to wear full protective suits and charcoal respirators. This is impractical and potentially dangerous in high summer temperatures. A deep placement coultter system (Avenger, Yetter Manufacturing Co, Colchester, IL) was modified to permit injection of 1,3-D into undisturbed soil at 30 cm depths and seal the soil above the injection point without creating channels for the fumigant to escape (John Mirusso, Mirusso Fumigation and Equipment, Delray Beach, FL). Ten large-scale field trials on a total of 31.5 ha were conducted using this technology during 2000–2001. Herbicide applications were combined with different formulations and rates of 1,3-D + chloropicrin and applied using the deep placement coultter system. In some trials, chloropicrin was shank injected into beds prior to laying plastic. In all ten trials, no difference in the density of plant parasitic nematodes was observed between experimental treatments and adjacent methyl bromide fields. Similar levels of weed control were observed in eight of ten trials, but the chemical alternatives failed to provide weed control in two trials conducted on farms with high populations of nightshade (*Solanum nigrum* L). Lack of control was attributed to grower inexperience with broadcast applications of pre-emergence herbicides. In the field trials where fumigant applications were made between November and February, levels of disease control similar to those with methyl bromide were achieved with a broadcast application of 1,3-D and chloropicrin. When fumigant applications were made between July and September, an additional application of chloropicrin in the bed was required to achieve levels of disease control similar to those with fumigation with methyl bromide, due to the increased disease pressure during the warmer crop production months.

Application methods and technology were modified to make soil solarization and broadcast fumigation compatible with Florida crop production systems. In both cases, adaptation to local production systems enhanced their performance. When used as a single tactic, soil solarization and 1,3-D + chloropicrin did not provide the same broad level of pest control as methyl bromide. A comprehensive pest management program that included additional tactics such as herbicides, fumigants and/or resistant cultivars was required to provide pest control similar to methyl bromide on a consistent basis.

3 CALIFORNIA STRAWBERRY PRODUCTION

3.1 Overview

California has the most productive strawberry fields in the world due to 50 years of research on optimizing cultivars and cropping practices in the context of soil fumigation with methyl bromide + chloropicrin (MBC). The production system is also based on vigorous, clean transplants, which are grown in MBC-fumigated nursery soils. In fruiting and nursery fields, fumigation with MBC is essential for weed,

disease and nematode control, and non-specific growth promotion. In fruiting fields, fumigation insures a return on the investment required for crop establishment ($\sim \$25\,000\text{ ha}^{-1}$) and total crop production ($\sim \$62\,000\text{ ha}^{-1}$).

ARS research has focused on alternatives that can be implemented immediately with little change in cultural practices. Large multidisciplinary projects test currently available chemicals in fruiting and nursery fields as one-for-one replacements.^{31,32} Tests of alternative chemicals, application technologies and the use of alternative tarps for fumigation have provided growers with hope of short-term sustainability.^{31,33–35} Movement and distribution of the fumigants are being evaluated to optimize the use of new chemicals.^{36,37} Plant pathologists have used many of the same chemical trials to document the influence of alternative chemicals on specific plant pathogens^{38,39} and potentially beneficial micro-organisms.^{39,40} Understanding the ecology of these organisms is important for controlling plant diseases in the absence of fumigation.

To avoid potential regulatory restrictions on chemicals under evaluation, non-chemical alternatives to chemical fumigants are being investigated. Rotations with broccoli reduced inoculum of the most important soil-borne pathogen of strawberry, *Verticillium dahliae* Kleb, and increased yield.⁴¹ Microbial inoculants are being evaluated and developed for soilborne diseases^{39,40,42} and will be used in conjunction with other management strategies for control of weeds.

ARS has taken the lead on a multi-agency and multidisciplinary (entomology, weed science, plant pathology and sociology) project, which enlists growers to help design, improve and implement a biologically integrated production system for strawberries. The new practices provided plant growth stimulation and weed control (Reference 43 and Fennimore S, pers comm) but improvement in the system is needed before adoption will occur.

3.2 Research highlight: organic strawberry production

ARS is also working to improve alternative cropping systems that are already in place. Despite the high risk involved in strawberry production in the absence of methyl bromide, the acreage of certified organic strawberry production has increased. The number of organic strawberry growers certified by a leading certification organization in California has increased by 27% from 34 to 46 from 1999 to 2001.^{44,45} Experiments comparing organic and conventional production have reported yields for the organic plots as high as 72% of conventional yields.⁴⁶ That growers are able to attain these yields, often on marginal land and with virtually no research support, suggests that research conducted in an organic context should help to optimize their production systems.

Although choice of variety is very important for success, the performance of commercially available

cultivars had never been compared in organic fields. High yielding cultivars were evaluated and selected for their yield and fruit quality in conventional production practices.⁴⁷ Organic growers rely on cultivars designed for use in MBC-fumigated soils, although cultivars perform differently in fumigated and non-fumigated conventional fields.⁴⁸ Because, in addition to the lack of fumigation, organic producers do not use fungicides or pesticides, it is likely that these differences in performance will be exaggerated in organic fields. This makes cultivar comparisons in conventional fields less relevant for cultivar selection for organic production.

A comparison of commercially available cultivars in organic production fields was completed by ARS scientists and collaborators. Experiments compared 15 cultivars at three locations during two field seasons in the central coast region of California. Overall, the cultivars Aromas, Pacific and Seascape performed the best in these trials.⁴⁹ Aromas, Seascape and Pacific have each been shown to be tolerant to at least one of the following soilborne pathogens: *Pythium ultimum* Trow, *Phytophthora cactorum* Schroet and *Verticillium dahliae* Kleb.^{38,39,50}

Planting material is about to become a more significant problem for organic growers. The USDA's new National Organic Standards will require that growers obtain a quality of transplant that is currently not available. Currently organic growers use conventionally grown strawberry transplants as a default because organically grown transplants are not available. Disease-free organic transplants can be produced under greenhouse conditions. In an organic production field, the performance of organic plug plants was compared with conventional field transplants. The experiment was a randomized complete block experiment with four replications. Berries were harvested once a week with cull and market quality fruit weighed separately. Data for yield are the cumulative data for all harvests. Conventional transplants were planted as recommended for adequate chilling. Plug plants were planted on the same day as bare-root transplants. There were significant differences between conventionally produced plants and organic plug plants for market ($P = 0.002$) and total ($P = 0.0009$) yield. Plug plants had significantly lower yields than the conventional transplants (Fig 1). Additional research is needed to determine optimum planting dates and conditions for production from organic plug plants. It will take time to optimize any new production system to achieve yields comparable with the current system, which has benefited from five decades of farmer innovation and scientific research.

4 FLORICULTURE AND ORNAMENTALS

The floriculture and ornamental industry is highly diverse and includes the production of ornamental nursery plants, potted plants, cut flowers, foliage and bulb crops. The California floriculture industry had a wholesale value of \$842 million in 2000.⁵¹




Figure 1. Market and total yield of conventional transplants and organic plug plants grown in a certified organic strawberry production field. Perpendicular lines at the top of the bars represent the standard error of the mean.

The cut flower industry in California was worth approximately \$286 million.⁵¹ In Florida, the value of this commodity is approximately \$29 million.⁵² Florida produces approximately 95% of the world's caladiums (*Caladium* spp),⁵³ which had a value of \$15 million last year (Terri Cantwell, Bates Sons and Daughters, pers comm). The diversity of crops that are grown in the floriculture and ornamentals nurseries in California and Florida represents a very complex research problem. While the total methyl bromide consumption by these commodities is difficult to estimate, the loss of this fumigant will present a unique challenge to these growers. The ornamentals industry is composed of hundreds of species of crops and thousands of varieties. Due to the number of different crops that are grown concurrently and in succession, the issue of phytotoxicity of alternative chemicals is extremely important. In addition to having severe limitations on the number of registered pesticides that are labeled for these crops, there is a requirement for clean propagation material in order for it to be shipped between states and to other countries. Weed control with a limited number of labeled herbicides and the need for clean, certifiable stock are the greatest concerns for growers of these commodities.

USDA research in this area is relatively new, although there are many technologies that have been developed for other crops that may have application for floriculture and ornamental production. Soil solarization has potential for these crops and some pathogens that have been found to be problems in ornamentals can be controlled in this manner.⁵⁴ Botanical extracts and essential oils have also been investigated as alternatives to methyl bromide for control of *Fusarium* wilt caused by *Fusarium oxysporum* Schlecht f sp *chrysanthemi* Luttr, GM Armstrong & JK Armstrong. Soil infested with this pathogen was treated with aqueous emulsions of formulated extracts of clove, neem, chili pepper extract and essential oil of mustard, and cassia. Ten per cent solutions of pepper/mustard, cassia, and clove extracts significantly reduced the soil populations of the pathogen.⁵⁵ These extracts could be used as a component in

a biologically based integrated pest management strategy for ornamentals.

5 PERENNIAL CROPS

Pre-plant methyl bromide use in orchards and vineyards in the USA totaled 2700 metric tons in 1997, accounting for approximately 15% of the total pre-plant use in the USA.⁵⁶ Perennial crops are not treated every cropping season, as are annual crops, but only when the orchard or vineyard is replanted. Grapes had a value of \$2.8 billion in California in 2000, peaches a value of \$251 million, almonds a value of \$681 million and walnuts a value of \$289 million.⁵⁷ 'Replant disorder' is a general term for the lack of vigor in newly replanted orchards and vineyards as compared with trees and vines planted in 'non-replant' soil. Fumigation with methyl bromide prior to replanting alleviates this problem. Fumigation with 1,3-D has also been used to control replant disorder, but California's township limits for 1,3-D restrict the availability of this option in some areas.⁵⁸ Perennial crops have deep root systems, some of which remain in the soil after old trees or vines are removed. These deep roots can serve as a reservoir of pathogens and nematodes, ready to infect the new trees and vines as soon as they are planted. Field trials to evaluate potential alternatives for perennial crops must determine efficacy of control, not only at the time of planting but also the on-going performance during the early growth and fruiting years. Thus, each field trial requires several years and significant resources before sufficient data are obtained.

5.1 Overview

ARS research has investigated biological, chemical, cultural and genetic approaches to finding alternatives to methyl bromide for perennial crops. Apple replant disorder has been characterized as largely fungal in nature. A crop of wheat prior to replanting apple fostered a disease-suppressive microbial community.⁵⁹ Vines planted in fumigated soil responded to inoculation with vesicular-arbuscular (VA) mycorrhizae, suggesting that the beneficial organisms that are suppressed by some chemical treatments are necessary for vigorous vine growth and should be reintroduced following fumigation.⁶⁰ Chloropicrin, 1,3-D and 1,3-D + chloropicrin have resulted in tree and vine growth as good or better than with methyl bromide.^{61–63} Shank-applied iodomethane gave good tree growth, but vine growth was intermediate between methyl bromide and untreated controls.^{62,64} Recent trials indicate that shank- and drip-applied iodomethane and propargyl bromide and drip-applied chloropicrin, azide and 1,3-D + chloropicrin each controlled plant parasitic nematodes as well as methyl bromide at planting, but long-term efficacy is not yet known.⁶⁵ Fallowing for 3 years resulted in tree growth as good as or better than with methyl bromide for plum, but not for peaches.⁶³ A 3-year fallow in a grape replant trial resulted in

a significant reduction in *Meloidogyne spp.*, rootknot nematode, but not in *Tylenchulus semipenetrans* Cobb, citrus nematode. Tree and vine rootstocks can be resistant to a specific pest, but susceptible to others.^{62,66} Accurate diagnosis of the problem in a specific orchard or vineyard will be critical to the selection of the best management option.

5.2 Research highlight: orchard replant

Replant disease of apple is typically controlled through the application of pre-plant soil fumigants, including methyl bromide, prior to orchard establishment on old orchard sites. Soil fumigation has been the option of choice due to the uncertain etiology of the disease. Systematic studies conducted in multiple orchards in Washington state utilized several approaches to define the causative agents of replant disease.⁶⁷ A fungal complex consisting of various species in the genera *Cylindrocarpon*, *Phytophthora*, *Pythium* and *Rhizoctonia* was shown to be the dominant cause of replant disease. Although implicated as a dominant causal agent in other geographical regions, the lesion nematode, *Pratylenchus penetrans* (Cobb) Filipjev & Stekhoven, was shown to have a limited site-specific role in disease development in Washington, a finding consistent with previous reports.⁶⁸ Identification of the causal pathogen complex has enabled the formulation of biologically based management options for the control of apple replant disease.

Based on composition of the causal pathogen complex, several biological, cultural and narrow-spectrum chemical control options have been devised and are being evaluated. The application of selected independent treatments has, in certain instances, yielded promising results. Physical manipulation of the orchard environment through altering spatial arrangements of the orchard or soil disturbance, minimized the impact of replant disease and enhanced yield of Gala/M.26 at the Columbia View orchard. Soil disturbance entailed excavation to a depth of 0.5 m in the fall prior to planting and spreading the soil over the adjacent ground. This resulted in soil exposure to repeated freeze/thaw cycles, an event that can limit the survival and activity of certain soil-borne pathogens, including *Rhizoctonia solani* Kühn.⁶⁹ Trees were also established in the old orchard aisle, rather than tree row, which subjected newly established trees to an environment possessing a reduced disease potential. Both treatments enhanced growth and yield of Gala/M.26 at this site (Table 2). As the pathogen complex at this site was comprised of *Cylindrocarpon destructans* (Zinssm) Scholten, *Phytophthora cactorum* (Lebert & Cohn) Schroeter, *Pythium heterothallicum* Campbell & Hendrix and *Rhizoctonia solani* AG 6, but not *Pratylenchus penetrans*,⁶⁷ a soil drench consisting of the fungicides metalaxyl and flutolanil was employed. Although vegetative growth was initially suppressed by this fungicide treatment, disease control and yield at the second harvest were comparable to that obtained through methyl bromide fumigation (Table 2).

Table 2. Effect of cultural, biological and chemical methods on yield of apples cv 'Gala'/M26 planted on replant ground in 1998 at Columbia View orchard, Orondo, WA

Treatment	Year 2000 yields (kg per tree) ^a	Year 2001 yields (kg per tree) ^a
Untreated Control	4.6	20.64
Methyl bromide fumigation	7.2*	27.12*
Soil excavation	5.4	25.72*
Interplanting (aisle)	6.4*	— ^b
<i>Pseudomonas putida</i> 2C8	4.1	21.36
RootShield® (<i>T. harzianum</i>)	4.7	22.45
Difenconazole	3.4	23.71
Metalaxyl+flutolanil	4.5	29.1*
Humic acid	3.4	19.9

^a Means in a column followed by (*) are significantly different ($P = 0.05$) from the control.

^b Trees removed from the aisle October 2000.

Table 3. Suppression of root infection by *Rhizoctonia solani* AG 5 and enhanced apple growth induced through wheat cultivation of WVC-Auvil orchard soil occurs in a cultivar specific manner

Treatment	Root wt (g) ^{ab}	Shoot wt (g) ^{ab}	% Root infection ^{ab}
Control	0.47 a	1.11 a	18.2 b
Pasteurized (95 C) ^c	0.89 bc	1.78 a	42.7 c
Hill-81	0.58 ab	2.59 b	16.7 b
Madsen	0.66 ab	2.42 b	18.9 b
Lewjain	1.04 c	4.79 c	3.3 a
Penawawa	0.94 bc	4.04 c	1.4 a

^a After the wheat cultivation treatment, soil was infested with oat-bran inoculum^{69,70} of *R. solani* AG 5 isolate 5–103 at a rate of 1 g kg⁻¹.

^b Means in the same column followed by the same letter are not significantly different ($P = 0.05$).

^c Soils were cultivated to Lewjain wheat prior to pasteurization.

Induction of soil suppression of elements of the causal pathogen complex may have potential as a component in an integrated system for control of apple replant disease. In greenhouse trials, cultivation of multiple orchard replant soils with wheat induced soil suppression of an introduced isolate of *Rhizoctonia solani* AG 5. This pathogen incites *Rhizoctonia* root rot of apple⁷⁰ and contributes to replant disease development. Suppression was mediated through the resident soil microflora as evidenced by the elimination of disease suppression when soils were pasteurized following wheat cultivation but prior to introduction of *R. solani* (Table 3). It is interesting that the induction of disease suppression was found to vary depending on the wheat genotype.⁷¹ The wheat genotype-specific nature of the response was associated with specific transformations in the resident soil microbial community. One element of this transformation was the consistent selection by certain wheat genotypes of a fluorescent pseudomonad community that exhibited a significantly higher capacity to suppress *in vitro* growth of *R. solani* AG 5 that was resident in the non-treated replant orchard soils (Table 4). Wheat varieties that did not induce disease suppression supported a

Table 4. *In vitro* suppression of *Rhizoctonia solani* AG 5 (zones of inhibition in mm) induced by fluorescent *Pseudomonas* spp obtained from the roots of Gala apple seedlings grown in WVC-Auvil orchard soils cultivated to wheat

Wheat cultivar	Average zone of inhibition (mm) ^a
No Wheat (Control)	0.71 a
Hill-81	0.91 ab
Lewjain	2.32 c
Madsen	0.60 a
Penawawa	1.65 bc

^a Means in the same column followed by the same letter are not significantly ($P = 0.05$) different.

fluorescent pseudomonad community that did not differ from the control in its ability to suppress fungal growth.

Based on results of these studies and associated field trials it is apparent that, with the development of appropriate guidelines, the integration of biologically sustainable management practices has significant potential as an alternative to pre-plant soil fumigation for control of apple replant disease.

5.3 Research highlight: vineyard replant

A 65-year-old 'Thompson Seedless' vineyard, located at the USDA Parlier, CA, research station was selected for a grape replant field trial. The treatments are described in Table 5. Each treatment was replicated five times in a randomized complete block design.

Table 5. Treatments applied to a vineyard replant field

Treatment 1	Untreated control
Treatment 2	1-Year fallow
Treatment 3	1-Year fallow plus a sorghum-sudangrass hybrid cover crop
Treatment 4	Shanked application of methyl bromide, 99.5% methyl bromide, 0.5% chloropicrin (448 kg ha ⁻¹), tarped (the treated control)
Treatment 5	Shanked application of iodomethane (448 kg ha ⁻¹), tarped
Treatment 6	Combination application of Telone II EC (327 litres ha ⁻¹ of 1,3-D) in 60 mm of water through a buried drip tape plus Vapam (243 litres ha ⁻¹ of 42% metam-sodium) through microsprinklers
Treatment 7	Combination application of Telone II EC (327 litres ha ⁻¹ of 1,3-D) in 60 mm of water through a buried drip tape plus Vapam (243 litres ha ⁻¹ of 42% metam-sodium) through microsprinklers + 1-year fallow
Treatment 8	Combination application of Telone II EC (327 litres ha ⁻¹ of 1,3-D) in 100 mm of water through a buried drip tape plus Vapam (243 litres ha ⁻¹ of 42% metam-sodium) through microsprinklers
Treatment 9	Combination application of Telone II EC (327 litres ha ⁻¹ of 1,3-D) in 100 mm of water through a buried drip tape plus Vapam (243 litres ha ⁻¹ of 42% metam-sodium) through microsprinklers +1-year fallow

Vines were removed from the 1-year fallow plots in fall, 1996. All other vines were removed in fall, 1997. 1,3-D + metam-sodium treatments were applied in early January, 1998. Methyl bromide and iodomethane treatments were applied in late April, 1998. In July of 1998, each plot was planted with three grape variety/rootstock combinations; own-rooted Thompson Seedless, Merlot on Harmony rootstock and Merlot on Teleki 5C rootstock. The rootstocks vary in levels of resistance to nematodes, which are thought to play a role in replant disorder. First- and second-year results of this study were reported previously.^{72,73}

Soil samples were collected to a depth of 60 cm from each treatment/rootstock combination in June, 2001 and processed by sugar flotation–centrifugation.⁷⁴ Although seven different plant parasitic nematode genera occurred in the field, the predominant genera were the root-knot (*Meloidogyne spp*) and the citrus (*T semipenetrans*) nematode. Population levels for these two genera are given in Table 6. Three years after treatment, the 1,3-D + metam-sodium combinations and iodomethane continue to give reductions of population levels of the root-knot and citrus nematode comparable with that of methyl bromide. While still present, the reduction in root-knot nematode populations observed in previous years in the 1-year-fallow and 1-year-fallow + cover-crop treatments, is no longer statistically significant in Thompson seedless and Harmony plots. Untreated

control and 1-year-fallow + cover-crop plots planted to the Harmony rootstock had significantly lower rootknot nematode populations than plots planted to Thompson seedless or Teleki 5C. Harmony supported the highest populations of citrus nematode and Teleki 5C the lowest. The difference was significant for both untreated and 1-year fallow plots.

Three years after treatment and re-planting, drip-applied 1,3-D and shank-applied iodomethane continue to give control of the root-knot and citrus nematode populations that is equivalent to that obtained with methyl bromide. The Harmony rootstock continues to support only minimal populations of the root-knot nematode, even in the untreated plots, but supports higher populations of the citrus nematode than either Thompson Seedless or Teleki 5C. Resistant rootstocks can be effective, but are more expensive than own-rooted vines and often not resistant to the diversity of pests that is encountered in a replant situation. If only root-knot nematode is present, the Harmony rootstock is a good alternative, alone or in combination with chemical controls, but citrus nematode populations will increase on Harmony if it is used without any chemical control. Iodomethane and the 1,3-D + metam-sodium combinations appear to be good alternatives to methyl bromide for vineyard replant when both root-knot and citrus nematode are present, at least for the first 3 years after establishment of a replanted vineyard.

Table 6. Nematode populations per 100 ml soil in soils receiving different treatments^{abc}

Treatment	<i>Meloidogyne sp</i>			<i>Tylenchulus semipenetrans</i>		
	Thompson Seedless	Teleki 5C	Harmony	Thompson Seedless	Teleki 5C	Harmony
Untreated control	103.7 a	71.7 a	1.0 a	972.8 a	334.1 a	1216.0 a
1-Year fallow	85.8 a	27.7 b	0.0 a	661.1 a	291.8 a	1562.9 a
1-Year fallow plus cover crop	50.7 a	16.2 b	2.2 a	573.4 a	275.2 a	921.6 a
Methyl bromide (448 kg ha ⁻¹)	2.9 b	0.0 c	0.0 a	0.5 bc	2.2 b	108.2 b
Iodomethane (448 kg ha ⁻¹)	7.8 b	0.0 c	0.0 a	12.2 bc	3.0 b	58.7 b
Telone II EC (327 litres ha ⁻¹ of 1,3-D applied in 60 mm of water) + Vapam (243 litres ha ⁻¹)	0.2 b	0.0 c	4.5 a	1.1 bc	0.5 b	49.9 b
Telone II EC (327 litres ha ⁻¹ of 1,3-D applied in 60 mm of water) + Vapam (243 litres ha ⁻¹) + 1-year fallow	0.2 b	0.0 c	0.0 a	0.0 c	0.0 b	5.6 b
Telone II EC (327 litres ha ⁻¹ of 1,3-D applied in 100 mm of water) + Vapam (243 litres ha ⁻¹)	33.9 b	0.0 c	0.0 a	37.4 b	15.4 b	4.5 b
Telone II EC (327 litres ha ⁻¹ of 1,3-D applied in 100 mm of water) + Vapam (243 litres ha ⁻¹) + 1-year fallow	21.8 b	0.0 c	0.0 a	7.5 bc	1.1 b	59.5 b

^a Mean of five replicates, soil sampled June 2001.

^b Drip treatments applied January 1998; shank treatments applied April 1998.

^c Means for each nematode genus/cultivar or nematode genus/rootstock combination followed by the same letter are not significantly different ($P = 0.05$).

5.4 Perennial field nursery crops

Soil fumigation with methyl bromide has commonly been used prior to planting field nurseries to insure that planting material grown for commercial and home-owner planting is free of soil-borne pathogens. Clean planting stock is critical not only for conventional growers, but also for organic growers. The California Code of Regulations makes it 'mandatory that nursery stock for farm planting be commercially clean with respect to economically important nematodes.'⁷⁵ The standard nursery treatment is methyl bromide. Fumigation with 1,3-D can be acceptable under some field conditions,⁷⁵ but its use is limited in California by township caps.⁵⁸ Growers of perennial nursery crops need an alternative to methyl bromide in order to continue to produce clean planting material and, if in California, to meet California Department of Food and Agriculture (CDFA) requirements. In order to be an acceptable management strategy, an alternative must be effective, available, economical and environmentally acceptable. In addition, to be approved as a nursery treatment in California, control of plant parasitic nematodes must be demonstrated to a soil depth of 152 cm (5 feet). In a grapevine nursery trial, shank-injected iodomethane and propargyl bromide and drip-applied chloropicrin, 1,3-D + chloropicrin, iodomethane, propargyl bromide and sodium azide were comparable with methyl bromide in the level of control of plant parasitic nematodes achieved to a depth of 152 cm.⁶⁵ Studies have been initiated recently to determine the impact of alternative fumigants and application technologies in garden rose and retail ornamental nurseries in California, but data are not yet available.

6 POST-HARVEST

Methyl bromide is used as a post-harvest fumigant to preserve product quality and to prevent the undesirable movement of insects, pathogens and nematodes that could be transported with commodities. Although quarantine use of methyl bromide is exempted under the Montreal Protocol, the loss of the larger pre-plant market for methyl bromide could result in reduced economic interest by manufacturers in continued production of this material. Loss of methyl bromide could mean the potential loss of several hundred million dollars in imports and exports of fruit, vegetables and nuts. ARS research programs have developed potential chemical and non-chemical replacement treatments for post-harvest use of methyl bromide.

6.1 Chemical alternatives

For fresh and durable commodities, time is often the most important factor in marketing the commodity. Consequently, replacements for methyl bromide can be categorized into those fumigants that require one day to several days and those which require only a few hours to be effective. Short treatment times are critical for fresh commodities because damage

to the commodity is more likely to occur during long exposures. In addition, the large volumes of commodities to be treated often require quick turn-around time in the fumigation chamber. With this in mind, ARS scientists have concentrated on developing shorter time applications.

6.1.1 Iodomethane

Iodomethane is a logical choice to replace methyl bromide, but it is not registered for use on commodities. In trials using stored-product insects, it has been shown to be as active or more active than methyl bromide. In specific tests, iodomethane was effective in controlling pests such as Indianmeal moth, *Plodia interpunctella* (Hübner), navel orangeworm, *Amyelois transitella* (Walker) and codling moth, *Cydia pomonella* L.^{76–80} One potential problem observed with iodomethane is its high rate of adsorption onto commodities.

6.1.2 Carbonyl sulfide

Carbonyl sulfide was developed in Australia as a grain fumigant and is undergoing registration there. It has good penetrating action and is toxic to a variety of stored product insects.⁷⁸ In disinfestation trials, carbonyl sulfide gave 100% mortality of diapausing codling moth larvae inside walnuts, when exposed for 24 h at a dosage of 40 mg litre⁻¹. The sorption of carbonyl sulfide on walnuts was less than that of either methyl bromide or iodomethane. Carbonyl sulfide is not an ideal quarantine fumigant because it requires lengthy treatment times to be effective against the Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann).⁸¹ One problem that was identified with carbonyl sulfide is that an odor often remains in the commodity following fumigation.⁷⁶ These objectionable odors may be caused by either wet commodity or contamination of the formulation with hydrogen sulfide.⁸²

6.1.3 Sulfuryl fluoride

Sulfuryl fluoride (Profume®), used for many years as a structural fumigant under the trade name of Vikane®, has the advantage of being registered, but does not yet have food tolerances established. Studies have shown that sulfuryl fluoride is as active or more active against the adult, pupal and larval life stages, but less active against the egg stage, of insects than with methyl bromide.⁷⁶ Of the fumigants tested as alternatives, this compound exhibits the least sorption during fumigation and is an excellent penetrating fumigant.^{76,83,84} When used on lemons and nectarines, sulfuryl fluoride caused phytotoxicity.⁸²

6.1.4 Ozone

Ozone has the advantage of having a GRAS (Generally Considered As Safe) designation that exempts it from registration requirements. It has been shown to be effective against stored-product pests but it requires several hours when used alone and is a poor

penetrant.⁸⁵ It is more effective when used with low levels of carbon dioxide, and vacuum may increase its toxicity or decrease the time required to obtain 100% mortality. It will require some special fumigation equipment because of its corrosive action.

6.2 Non-chemical methods

Efforts to develop non-chemical treatments for post-harvest use have concentrated on inert gases, heat/cold treatments, and radio frequency (RF) energy. In addition, some non-direct methods such as trapping and mating disruption of insects in the field might preclude conventional post-harvest treatments of codling moth and navel orangeworm.^{86,87}

6.2.1 Heat/cold treatments

Hot air has proved a successful treatment for apple maggot, *Rhagoletis pomonella* Walsh.^{88,89} Furthermore, it can eliminate fruit flies from Valencia oranges,^{90,91} although some loss in flavor might occur due to loss in volatiles from the oranges.⁹² Neven and Mitcham⁹³ demonstrated that combining controlled atmospheres with hot air greatly reduces the treatment time when compared with hot air alone. Low temperatures have been investigated for the elimination of Indianmeal moth and navel orangeworm from dried fruit,⁹⁴ but the long treatment times (days) suggest that these cold treatments would best be used as control disinfestations rather than as quarantine treatments.

6.2.2 RF energy

Results have shown that using long-wavelength energy to heat cherries in saline solution can be used as a quarantine treatment.⁹⁵ Also research shows that RF energy treatments are effective against codling moth and navel orange worm in walnuts.^{96–98} At the wavelengths being studied, insects are selectively heated without adversely heating the fruit which might cause damage.

6.2.3 Irradiation

Irradiation has been shown to be the most efficient and least damaging method to treat some exposed fruit.^{99,100} Generally, doses from 50 to 200 Gy are sufficient for quarantine security, but the exact dose varies with the insect being targeted. Four basic problems exist for using irradiation: (1) the capital costs for the facility are large and require that it operate year-round, (2) large volumes of commodities must be treated very quickly to allow timely movement through the marketing channels, (3) irradiation can render the adult stages sterile, rather than dead, leaving the inspector uncertain as to whether the insects were exposed to irradiation, whether all the insects were treated, or whether the adult entered the commodity following irradiation, and (4) many countries will not accept irradiation as a commodity treatment.

6.2.4 Trapping/recycling of methyl bromide

In 1995 to 1998, a cooperative agreement between ARS and industry investigated the most efficient

method to capture methyl bromide on activated carbon. Studies showed that up to an 18% load of methyl bromide could be put on the activated carbon, depending upon temperature.¹⁰¹ This research led to a commercial product and now two adsorption units are installed on working fumigation chambers in the USA.

7 CONCLUSION

The ARS research program on alternatives to pre-plant and post-harvest use of methyl bromide covers a diversity of cropping systems and fresh and durable commodities, utilizing a range of approaches including biological, genetic, cultural, chemical and physical control strategies. The program encompasses both short-term objectives to help growers and industry cope in the near term with the impending ban, and long-term objectives to transition to more integrated, sustainable management systems. Several short-term alternatives have been identified and evaluated for both pre-plant and post-harvest uses. Some will require regulatory approval before they will be available for commercial use. Good progress is being made in component research programs that will serve as the foundation for the development of integrated systems that begin with pre-plant preparation and end with commodity treatments.

REFERENCES

- 1 FDACS, *Florida agricultural statistics, vegetable summary*, Florida Department of Agriculture and Consumer Services, Florida Agricultural Statistics Service. Orlando, FL, USA 63 pp (1999).
- 2 USDA, *Agricultural statistics*, National Agricultural Statistics Service (1997), available at <http://www.nass.usda.gov>
- 3 USDA, *Agricultural chemical usage vegetables*, National Agricultural Statistics Service (1999), available at <http://www.nass.usda.gov/fl/>
- 4 EPA, *Methyl bromide use*, United States Environmental Protection Agency (1997), available at <http://www.epa.gov/spdpublic/mbr>
- 5 Spreen TH, VanSickle JJ, Moseley AE, Deepak MS and Mathers L, Use of methyl bromide and the economic impact of its proposed ban on the Florida fresh fruit and vegetable industry, *Univ Fla Inst Food Agric Sci Bull*, No 898 (1995).
- 6 Geraldson CM, Pepper production efficiency using the gradient-mulch concept. *Proc Fla State Hort Soc* 90:152–155 (1977).
- 7 Geraldson CM, Overman AJ and Jones JP, Combination of high analysis fertilizers, plastic mulch, and fumigation for tomato production on old agricultural land. *Soil Crop Sci Soc Fla* 25:18–24 (1965).
- 8 Fery RL, Dukes PD and Ogle WL, 'Carolina Cayenne' Pepper. *HortScience* 21:330 (1986).
- 9 Thies JA and Mueller JD, Use of resistant pepper as a rotational crop to manage southern root-knot nematode. *HortScience* 33:716–718 (1998).
- 10 Kokalis-Burelle N, Vavrina CS, Rosskopf EN and Shelby RA, Field evaluation of PGPR amended transplant mixes and soil solarization for tomato and pepper production in Florida. *Plant and Soil* 238:257–266. (2001).
- 11 Kadir JB, Charudattan R, Stall WM and Brecke BJ, Field efficacy of *Dactylaria higginsii* as a bioherbicide for the control of purple nutsedge (*Cyperus rotundus*). *Weed Technol* 14:1–6 (2000).

- 12 Roskopf EN, Chellemi DO, Charudattan R and Chandramohan S, Alternatives to methyl bromide for weed control. *Acta Hort* 532:103–107 (2000).
- 13 Roskopf EN, Charudattan R, DeValerio JT and Stall WM, Field evaluation of *Phomopsis amaranthicola*, a biocontrol agent for *Amaranthus* spp. *Plant Dis* 84:1225–1230 (2000).
- 14 Larkin RP and Fravel DR, Efficacy of various fungal and bacterial biocontrol organisms for control of Fusarium wilt of tomato. *Plant Dis* 82:1022–1029 (1998).
- 15 Larkin RP and Fravel DR, Mechanisms of action and dose-response relationships governing biological control of Fusarium wilt of tomato by nonpathogenic Fusarium spp. *Phytopathology* 89:1152–1161 (1999).
- 16 Locascio SJ, Dickson DW and Roskopf EN, Alternative fumigants applied with PE and VIF mulches for tomato, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 17–1 to 17–3 (2001).
- 17 Noling JW, Gilreath JP and Roskopf EN, Alternatives to methyl bromide field research efforts for nematode control in Florida, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 14–1 to 14–3 (2001).
- 18 Roskopf EN, Albano JP and Lamb EM, ‘Green Chemistry’ for bell peppers: management of *Phytophthora capsici*, *Proc Fla Ag Conference and Trade Show*, 25 (2001).
- 19 Kokalis-Burelle N and Fuentes-Borquez P, Efficacy of PLANTPRO 45™ as an alternative to methyl bromide for control of root-knot nematode on tomato. *J Nematol* 32:440 (2000).
- 20 Adams PD and Kokalis-Burelle N, Efficacy of PLANTPRO 45™ as a seed treatment for the control of Fusarium wilt of basil. *Phytopathology* 90:S2 (2000).
- 21 Adams PD and Kokalis-Burelle N, Herbicidal effect of Plantpro45™ on germination of *Portulaca oleracea*. *Phytopathology* 91:S2 (2001).
- 22 Chellemi DO, Olson SM and Mitchell DJ, Effects of soil solarization and fumigation on survival of soilborne pathogens of tomato in northern Florida. *Plant Dis* 78:1167–1172 (1994).
- 23 McSorley R and Parrado JL, Application of soil solarization to rockdale soils in a subtropical environment. *Nematropica* 16:125–140 (1986).
- 24 Overman AJ, Off-season land management, soil solarization and fumigation for tomato. *Proc Soil Crop Sci Soc Fla* 44:35–39 (1985).
- 25 Chellemi DO, Olson SM, Mitchell DJ, Secker I and McSorley R, Adaptation of soil solarization to the integrated management of soilborne pests of tomato under humid conditions. *Phytopathology* 87:250–258 (1997).
- 26 Chellemi DO, McSorley R, Rich FR and Olson SM, Field validation of soil solarization for fall production of tomato. *Proc Fla State Hort Soc* 110:330–332 (1997).
- 27 Chellemi DO, Hochmuth RC, Winsberg T, Guetler W, Shuler KD, Datnoff LE, Kaplan DT, McSorley R, Dunn RA and Olson SM, Application of soil solarization to fall production of cucurbits and pepper. *Proc Fla State Hort Soc* 110:333–336 (1997).
- 28 Gilreath JP, Noling JW, Gilreath PR and Jones JP, Field validation of 1,3-dichloropropene + chloropicrin and pebulate as an alternative to Methyl Bromide in tomato. *Proc Fla State Hort Soc* 110:273–276 (1997).
- 29 Locascio SJ, Gilreath JP, Dickson DW, Kucharek TA, Jones JP and Noling JW, Fumigant alternatives to methyl bromide for polyethylene mulched tomato. *HortScience* 32:1208–1211 (1997).
- 30 Dow Elanco, Telone C-35 specimen and supplemental label for Florida, Dow Elanco Chemical Corp, USA (1998).
- 31 Ajwa H and Trout T, Strawberry growth and yield with three years of drip fumigation, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, Orlando, FL, USA, pp 25–1 to 25–4 (2000).
- 32 Fennimore S, Martin F, Duniway J, Browne G, Ajwa H, Winterbottom C, Westerdahl B, Goodhue R, Haar M and Bull C, Alternative fumigants for control of soil pests: Strawberry as a model system, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, p 96–1, (2001).
- 33 Ajwa H, Schutter M, Nelson S, Trout T and Winterbottom C, Efficacious application rates of propargyl bromide and iodomethane/chloropicrin for strawberry production, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 24–1 to 24–3 (2001).
- 34 Eayre CG and Sims JJ, Effect of methyl iodide and chloropicrin on strawberry yield in California. *Phytopathology* 90:S21 (2000).
- 35 Trout T, USDA program to evaluate Propargyl Bromide as a soil preplant fumigant, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 23–1 to 23–2 (2001).
- 36 Ajwa H, Trout T, Nelson S and Schutter M, Drip fumigation: Water and fumigant distribution in soil, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 50–1 to 50–2 (2001).
- 37 Yates SR, Dugan R and Papiernik SK, Predicting pathogen control from soil fumigation, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 19–1 to 19–3 (2001).
- 38 Browne GT, Becherer HE, Vazquez MR, McGlaughlin SA, Wakeman RJ, Winterbottom CQ, Duniway JM and Fennimore SA, Outlook for managing Phytophthora diseases on California strawberries without methyl bromide, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 29–1 to 29–2 (2001).
- 39 Martin FN, Management of pathogens associated with black root rot of strawberry, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 46–1 to 46–4 (2001).
- 40 Bull CT, Shetty KG and Subbarao KV, Interactions between myxobacteria, plant pathogenic fungi and biocontrol agents, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, Orlando, FL, USA, p 94–1 (2000).
- 41 Subbarao KV, Martin F and Koike S, Crop rotation with broccoli for management of Verticillium wilt, *The Pink Sheet (California Strawberry Commission Newsletter)* pp 1–11 (2001).
- 42 Eayre CG, Chloropicrin fumigation followed by bacterial treatment increases strawberry yield, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 100–1 to 100–2 (2001).
- 43 Bull CT, Participatory research for the dynamic California strawberry industry, *Proc Western Region SARE Conference*, Portland, OR, USA (March 2000).
- 44 California Certified Organic Farmers, CCOF Certified Organic Membership Directory, Santa Cruz, CA, USA (1999).
- 45 California Certified Organic Farmers, CCOF Certified Organic Membership Directory, Santa Cruz, CA, USA (2001).
- 46 Gliessman ST, Werner MR, Allison J and Cochran J, A comparison of strawberry plant development and yield under organic and conventional management on the central California coast. *Biol Agric Hort* 12:327–338 (1996).
- 47 Larson KD and Shaw DV, Relative performance of strawberry genotypes on fumigated and nonfumigated soils. *J Amer Soc Hort Sci* 120:274–277 (1995).
- 48 Martin FN, Management of root diseases of strawberry. *Phytopathology* 91:S58 (2001).
- 49 Bull CT, Koike ST and Shennan C, Performance of commercially available strawberry cultivars in organic production fields, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 28–1 to 28–4 (2001).
- 50 Duniway JM, Hao JJ, Dopkins DM, Ajwa H and Browne GT, Chemical, cultural, and biological alternatives to methyl bromide for strawberry, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 41–1 to 41–3 (2001).

- 51 California Agricultural Statistics Service, *California Floriculture*, (2000), available at <http://www.nass.usda.gov/cal/>
- 52 Florida Agricultural Statistics Service, *Foliage, Floriculture and Cut Greens*, (2001), available at <http://www.nass.usda.gov/fl/>
- 53 Sweet K, The Caladium crossing, *Ornamental Outlook*, March (1998).
- 54 Coelho L, Mitchell DJ and Chellemi DO, Thermal inactivation of *Phytophthora nicotianae*. *Phytopathology* **90**:1089–1097 (2000).
- 55 Bowers JH and Locke JC, Effect of botanical extracts on the population density of *Fusarium oxysporum* in soil and control of Fusarium wilt in the greenhouse. *Plant Dis* **84**:300–305 (2000).
- 56 National Center for Food and Agricultural Policy, *1997 Pesticide Use in Crop Production by Crop and State* (2001), available at www.ncfap.org
- 57 California Department of Food and Agriculture, *Resource Directory 2001*, California Department of Food and Agriculture, State of California, 175 pp (2001).
- 58 California Dept of Pesticide Regulation, Suggested permit conditions for using 1,3-dichloropropene pesticides (fumigant), Enforcement Letter ENF 01–40 to County Agricultural Commissioners, Aug 7, 2001.
- 59 Mazzola M, Managing soil microbial communities to enhance growth of apple in replant soils, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 43–1 to 43–4 (1999).
- 60 Linderman RG and Davis EA, Comparative response of selected grapevine rootstocks and cultivars to inoculation with different mycorrhizal fungi. *Am J Enol Vitic* **52**:8–11 (2001).
- 61 Browne G, Bulluck III LR, Connell J, Trout T and Schneider S, Determining unknown causes of replant disorder of *Prunus* species in California, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 102–1 to 102–4 (2001).
- 62 Schneider SM, Ajwa H, Trout T, Browne G and Sims J, On-going performance of vineyard replant trials initiated in 1998–2000, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 75–1 to 75–5 (2001).
- 63 Trout T, Ajwa H and Schneider S, Fumigation and fallowing effects on replant problems in California peach, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 77–1 to 76–7 (2001).
- 64 Eayre CG, Sims JJ, Ohr HD and Mackey B, Evaluation of methyl iodide for control of peach replant disorder. *Plant Dis* **84**:1177–1179 (2001).
- 65 Schneider SM, Ajwa H, Trout T, Browne G and Sims J, Field evaluation of nematode control alternatives for grapevine nurseries, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 76–1 to 76–4 (2001).
- 66 Nyczepir AP, Evaluation of peach rootstocks for management of root-lesion nematode in the Southeast, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, Orlando, FL, USA, pp 15–1 to 15–2 (2000).
- 67 Mazzola M, Elucidation of the microbial complex having a causal role in the development of apple replant disease in Washington. *Phytopathology* **88**:930–938 (1998).
- 68 Covey RP, Benson NR and Haglund WA, Effect of soil fumigation on the apple replant disease in Washington. *Phytopathology* **69**:684–686 (1979).
- 69 Mazzola M, Johnson TE and Cook RJ, Influence of field burning and soil treatments on growth of wheat after Kentucky bluegrass, and effect of *Rhizoctonia cerealis* on bluegrass emergence and growth. *Plant Pathol* **46**:708–715 (1997).
- 70 Mazzola M, Identification and pathogenicity of *Rhizoctonia* spp isolated from apple roots and orchard soils. *Phytopathology* **87**:582–587 (1997).
- 71 Mazzola M and Gu YH, Wheat genotype-specific induction of soil microbial communities suppressive to disease incited by *Rhizoctonia solani* anastomosis group (AG)-5 and AG-8. *Phytopathology* **92**:1300–1307 (2002).
- 72 Schneider SM, Ajwa H, Trout T and Sims J, Grape replant disorder—field tests of some potential alternatives, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 50–1 to 50–4 (1999).
- 73 Schneider SM, Ajwa H, Trout T and Sims J, Alternatives for vineyard replant and grapevine nurseries, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, Orlando, FL, USA, pp 14–1 to 14–5 (2000).
- 74 Jenkins WR, A rapid centrifugal-flotation technique for separating nematodes from soil. *Pl Dis Reporter* **48**:692 (1964).
- 75 California Dept of Food and Agriculture, Approved treatment and handling procedures to ensure against nematode pest infestation of nursery stock, *Nursery Inspection Procedures Manual*, Item #12, 18 pp (1996).
- 76 Zettler JL, Leesch JG and Gill RF, Chemical treatment alternatives for postharvest pests, 24th Annual Meeting Workshop on Phytosanitary Alternatives to Methyl Bromide. San Diego, CA, October 19, 2000, *NAPPO Bull*, No 16, pp 32–35, May (2001).
- 77 Zettler JL, Leesch JG, Gill RF and Tebbets JC, Chemical alternatives for methyl bromide and phosphine treatments for dried fruits and nuts, in *Proceedings of the Seventh International Working Conference on Stored Product Protection*, 1998, October 14–19, Beijing, China, ed by Jin Z, Liang Y, Tan X and Guan L, Chengdu (China), Sichuan Publishing House of Science and Technology, pp 554–561 (1999).
- 78 Zettler JL, Leesch JG, Gill RF and Mackey BE, Toxicity of carbonyl sulfide to stored product insects. *J Econ Entomol* **90**:832–836 (1997).
- 79 Leesch JG and Zettler JL, The fumigation of walnuts with carbonyl sulfide or methyl iodide as replacements for methyl bromide to control stored product pests, in *Walnut research reports 1998*, Walnut Marketing Board, Sacramento, CA, USA, pp 257–265 (1999).
- 80 Leesch JG and Zettler JL, The effectiveness of fumigating walnuts with carbonyl sulfide, methyl iodide, or sulfur dioxide in controlling stored product pests, in *Walnut research reports 1999*, Walnut Marketing Board, Sacramento, CA, USA, pp 227–232 (2000).
- 81 Obenland DM, Jang EB, Aung LH and Zettler JL, Tolerance of lemons and the Mediterranean fruit fly to carbonyl sulfide quarantine fumigation. *Crop Protect* **17**:219–224 (1998).
- 82 Aung LH, Leesch JG, Jenner JF and Grafton-Cardwell EE, Effects of carbonyl sulfide, methyl iodide, and sulfur dioxide on fruit phytotoxicity and insect mortality. *Ann Appl Biol* **139**:93–100 (2001).
- 83 Bell CH and Savvidou N, The toxicity of Vikane (sulfuryl fluoride) to age groups of eggs of the Mediterranean flour moth (*Ephestia kuehniella*). *J Stored Prod Res* **35**:233–247 (1999).
- 84 Xu G, Cheng Z, Seng Z and Qui N, The development of sulphuryl fluoride (SO₂F₂) in China—a brief introduction, in *Proceedings of the Seventh International Working Conference on Stored Product Protection*, 1998, October 14–19, Beijing, China, ed by Jin Z, Liang Y, Tan X and Guan L, Chengdu (China), Sichuan Publishing House of Science and Technology, pp 562–566 (1999).
- 85 Leesch JG, The mortality of stored-product insects following exposure to gaseous ozone at high concentrations, *Proceedings of the Annual Meeting of the International Ozone Association: Pan American Group, Advances in Ozone Technology*, Newport Beach, CA, USA, p 41 May 5–9, (2001).
- 86 Burks CS, Gerber RG and Brandl DG, Mating disruption for control of the navel orangeworm in the orchard, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 61–1 to 61–3 (2001).
- 87 Kuenen LPS and Rowe HC, Attractants for navel orangeworm to reduce infestations and aflatoxins, *Proc Annu Intl Res Conf*

- on Methyl Bromide Alternatives and Emissions Reductions, San Diego, CA, USA, p 111-1 (2001).
- 88 Hallman GJ, Potential quarantine treatments against plum curculio to replace methyl bromide, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, Orlando, FL, USA, pp 121-1 to 121-2 (1998).
- 89 Blessington TR and Hallman GJ, Hot-air quarantine treatment for apples infested with apple maggot. *Subtropical Plant Sci* 51:1-3 (1999).
- 90 Shellie KC and Mangan RL, Postharvest quality of Valencia orange after exposure to hot, moist forced air for fruit fly disinfestation. *HortScience* 29:1524-1527 (1994).
- 91 Obenland DM, Arpaia ML and Aung LH, Quality of nectarines subjected to forced-air heat treatment for Mediterranean fruit fly disinfestation. *J Hort Sci Biotech* 74:553-555 (1999).
- 92 Obenland DM, Arpaia ML, Austin RK and Mackey BE, High-temperature forced-air treatment alters the quantity of flavor-based volatile constituents present in navel and Valencia oranges. *J Agric Food Chem* 47:5184-5188 (1999).
- 93 Neven LG and Mitcham EJ, CATTS (controlled atmosphere/temperature system): a novel tool for the development of quarantine treatments. *Amer Entom Spring*, 56-59 (1996).
- 94 Johnson JA and Valero KA, Response of navel orangeworm and indianmeal moth eggs to low temperature storage, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, San Diego, CA, USA, pp 65-1 to 65-2 (1999).
- 95 Ikediala JN, Tang J, Hansen JD, Drake SR and Wang S, Development of saline-water-immersion technique with RF energy as a postharvest treatment against codling moth in cherries. *Postharvest Bio Technol* 24:209-221. (2002).
- 96 Hanson J, Tang J, Mitcham E, Johnson J, Wang S and Ikediala J, Application of radio frequency treatments to control insects in walnuts, in *Walnut research reports 2000*, Walnut Marketing Board, Sacramento, CA, pp 297-300 (2001).
- 97 Wang S, Ikediala JN, Tang J, Hansen JD, Mitcham E, Mao R and Swanson B, Radio frequency treatments to control codling moth in in-shell walnuts. *Postharv Biol Technol* 22:29-38 (2001).
- 98 Wang S, Ikediala JN, Tang J and Hansen JD, Thermal death kinetics and heating rate effects for fifth-instar codling moth (*Cydia pomonella* L.). *J Stored Prod Res* 38:441-453. (2002).
- 99 Hallman GJ, Radiation quarantine treatment for blueberries to replace methyl bromide, *Proc Annu Intl Res Conf on Methyl Bromide Alternatives and Emissions Reductions*, Orlando, FL, USA, pp 80-1 to 80-4 (1998).
- 100 Hallman GJ and Thomas DB, Gamma radiation quarantine treatment against blueberry maggot and apple maggot (Diptera: Tephritidae). *J Econ Entomol* 92:1373-1376 (1999).
- 101 Leesch JG, Knapp GF and Mackey BE, Methyl Bromide adsorption on activated carbon to control emissions from commodity fumigations. *J Stored Prod Res* 36:65-74 (2000).